


South Central US Regional Turf Spray Program

Bermudagrass and Zoysiagrass – Warm Season

For tees, greens or fairways, superintendents trust BASF for the most effective high-quality golf solutions. Proven BASF innovations deliver superior performance in all geographical regions and climates, especially during peak stress times. With a specific protocol for each area of the country, the BASF Regional Turf Spray Program is specially designed to provide proven disease control and produce stronger roots, greater stress tolerance, higher resiliency and increased growth efficiency. The result is championship conditions, all year long.

 **BASF**
We create chemistry



South Central US Bermudagrass and Zoysiagrass

Arkansas, Louisiana, Oklahoma and Texas

Bermudagrass greens go dormant in most areas of this region with very cold and icy conditions for some of the winter. Extended wet weather can also dominate the winter months, while warm conditions can develop slowly. This region can vary widely from hot and very dry, to cloudy and wet. Humidity increases farther south, and can last through the summer and into the fall.

21-28
day program



Timing	Product	Active Ingredient	Rate/1,000 sq. ft.	Key Target Diseases
1 Jan	Xzemplar ® fungicide + Mancozeb	Fluxapyroxad + Mancozeb	0.26 fl oz + SLR ¹ (Standard Local Rate)	Leaf spot ⁴ , algae
2 Feb	Segway + Mancozeb	Cyazofamid + Mancozeb	SLR ¹	<i>Pythium</i> , Leaf spot
3 Mar	■ Lexicon ® Intrinsic ® brand fungicide	Fluxapyroxad and Pyraclostrobin	0.34-0.47 fl oz ²	Bermudagrass decline, Take-all patch, Leaf spot, Fairy ring, Zoysia patch/Large patch ³
4 Apr	26GT + Signature Xtra Stressgard + Mancozeb	Iprodione + Fosetyl-AL (Aluminum tris) + Mancozeb	SLR ¹	Bermudagrass decline, Take-all patch, Leaf spot, <i>Pythium</i>
5 Apr	Prostar or Mirage Stressgard	Flutolanil or Tebuconazole	SLR ¹	Fairy ring, Zoysia patch/Large patch ³
6 May	Honor ® Intrinsic brand fungicide	Pyraclostrobin and Boscalid	1.11 oz	'Quickness to Closure-Healing' aerification, Leaf spot, Fairy ring, Zoysia patch/Large patch ³
7 Jun	Segway + Mancozeb	Cyazofamid + Mancozeb	SLR ¹	<i>Pythium</i> , Leaf spot
8 Jul	■ Lexicon Intrinsic brand fungicide	Fluxapyroxad and Pyraclostrobin	0.34-0.47 fl oz ²	Summer disease protection including Mini ring, 'Quickness to Closure-Healing' aerification
9 Jul	Prostar	Flutolanil	SLR ¹	Mini ring, Fairy ring
10 Aug	26GT + Mancozeb	Iprodione + Mancozeb	SLR ¹	Leaf spot
11 Sept	■ Lexicon Intrinsic brand fungicide	Fluxapyroxad and Pyraclostrobin	0.34-0.47 fl oz ²	Spring dead spot ⁵ (for those OK-AR areas), Leaf spot, <i>Pythium</i> , Zoysia patch/Large patch ³
12 Sept	Segway + Daconil	Cyazofamid + Chlorothalonil	SLR ¹	<i>Pythium</i> , Leaf spot, Algae
13 Oct	Xzemplar fungicide	Fluxapyroxad	0.26 fl oz	Fairy ring ⁴
14 Oct	■ Lexicon Intrinsic brand fungicide	Fluxapyroxad and Pyraclostrobin	0.34-0.47 fl oz ²	Spring dead spot ⁵ (for those OK-AR areas), Bermudagrass decline, Leaf spot, Fairy ring, Zoysia patch/Large patch ³
15 Nov	■ Honor Intrinsic brand fungicide	Pyraclostrobin and Boscalid	1.11 oz	Bermudagrass decline, Fairy ring
16 Dec	Signature Xtra Stressgard + Daconil	Fosetyl-AL (Aluminum tris) + Chlorothalonil	SLR ¹	<i>Pythium</i> , Leaf spot, Algae

■ *Holiday Spray Program*

¹ SLR = Standard Local Rate for the products shown; consult local extension recommendations in your region.

² **Lexicon Intrinsic** brand fungicide can be applied 3-4 times during any calendar year not to exceed 1.7 fl oz per 1,000 sq ft (74 fl oz per acre).

³ In those areas where there are zoysiagrass fairways, greens or tees, control fairy ring and zoysia patch/large patch caused by *Rhizoctonia solani* with applications in the fall, and repeat in the spring.

⁴ Subject to a BASF Recommendation under FIFRA 2(ee) for fairy ring, *Bipolaris* leaf spot and *Rhizoctonia* leaf spot and leaf and sheath spot in certain states. See www.CDMS.net for BASF Technical Information Bulletin.

⁵ Subject to a BASF Recommendation under FIFRA 2(ee) for spring dead spot in certain states. See www.CDMS.net for BASF Technical Information Bulletin.

Product timing is a suggestion only; optimize application for local weather and field conditions. Confirm active status of product registration in your state before using; not all products are registered in all states. Application Rate/1,000 sq. ft. is based on maximum label rate or range. Claims related to root strength, stress tolerance, resiliency and growth efficiency are specific to those BASF products containing pyraclostrobin, as shown.

Always read and follow label directions.

Honor, Intrinsic, Lexicon and Xzemplar are registered trademarks of BASF. 26GT, Mirage, Prostar, Signature and Stressgard are registered trademarks of Bayer. Segway is a registered trademark of ISK, Ltd. Daconil is a registered trademark of Syngenta Professional Products.

© 2017 BASF Corporation. All Rights Reserved.

RGSP-S19-01/17