







**■ BASF**

We create chemistry

## **Tower<sup>®</sup> herbicide**

### **Adaptive preemergent control**

Tower herbicide delivers selective preemergent control of annual grasses, sedges and small-seeded broadleaf weeds in golf course turf, lawns and ornamentals.

Challenging Weeds	Adaptive Utility	Premium Tank-Mix Partner	Enhanced Poa Defense
 <p>Preemergent control of goosegrass, doveweed, sedges, spurge, <i>Poa annua</i> and more</p>	 <p>Flexible rate range for warm-season and cool-season turf and short reseeding intervals</p>	 <p>Readily mixes with preemergent and postemergent herbicides to expand spectrum of control</p>	 <p>Residual control and early postemergent activity on <i>Poa annua</i></p>


### Advantages of Tower herbicide

- Selective preemergent control of annual grassy weeds, broadleaf weeds and sedges
- Inhibits shoot growth to control weeds as they germinate without root pruning effects on desirable turf
- Non-staining, liquid formulation readily mixes with other herbicides to expand spectrum of control
- Reliable activation with just 1/4" irrigation or rain
- Can be impregnated or coated onto dry bulk granular fertilizers

### Adaptive residual protection

- Flexible rate range allows for applications to actively-growing warm- and cool-season turfgrass
- Over-the-top tolerance on 100+ ornamental plant species in landscape beds
- Tank-mix compatible with postemergent herbicides for extended residual control on bareground and gravel sites

### Goosegrass residual control



### Turf best use recommendations

- Apply to dense and well-established turfgrass
- Make applications spring through fall when soil temperatures are above 55°F
- Use 21 fl oz / A on cool-season turf types (bentgrass, Kentucky bluegrass, fine fescue, perennial ryegrass and tall fescue)
- Use 21 to 32 fl oz / A on warm-season turf types (bahiagrass, bermudagrass, buffalograss, centipedegrass, kikuyugrass, St. Augustinegrass, seashore paspalum and zoysiagrass)
- Water-in application to rinse residue from foliage and activate herbicide barrier
- Make sequential applications 5 to 8 weeks apart

See product label for full use instructions

HRAC Group	Active Ingredients (%)	Usage Rate	Formulation	Packaging	Key Weeds Controlled
15	Dimethenamid-P (63.9%)	21 to 32 fl oz	Emulsifiable concentrate (EC)	2 x 2.5 gal	Goosegrass, doveweed, annual sedges, spurges, annual bluegrass, Florida pusley, eclipta, <i>Poa annua</i>

Always read and follow label directions.

[betterturf.basf.us](https://betterturf.basf.us)

Products may not be registered for sale or use in all states. Please check with your state or local Extension Service. Refer to label for registered uses. It is a violation of Federal law to use a product in a manner inconsistent with its labeling.

Tower is a registered trademark of BASF. Ronstar is a registered trademark of Environmental Science.  
©2024 BASF Corporation. All rights reserved. PSS 24-30-423